

AGASTIYA M-45

MANCOZEB 75% W.P. (CONTACT FUNGICIDE)

Caution: "Not To Be Used On Crops Other Than Specified On This Label/leaflet".
It is a co-ordinated product of Zinc & Manganese Enhance the fungicidal, bactericidal and contact fungicide activity of Mancozeb 75% WP, containing 75% WW active ingredient and balance adjuvants.

Common name of Crop/ Diseases	A.I. (Kg)	Doseage / Ha Formulation (Kg)	Dilution In Water (Lit)	Waiting Period last spray to harvest (In days)
Tomato	1.25-1.5	1.5-2	750	-
Early Blight	1.25-1.5	1.5-2	750	-
Late Blight	1.25-1.5	1.5-2	750	-
Buck Eye Rot	1.25-1.5	1.5-2	750	-
Leaf Spot	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Wheat	1.25-1.5	1.5-2	750	-
Black Rust	1.25-1.5	1.5-2	750	-
Blight	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Downy Mildew	1.25-1.5	1.5-2	750	-
Paddy	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Tip Rot	1.25-1.5	1.5-2	1000	-
Banana	1.25-1.5	1.5-2	1000	-
Sigatoka	1.25-1.5	1.5-2	1000	-
Leaf spot	1.25-1.5	1.5-2	1000	-
Cucumber	22.5 Gram/litre	30 Gram/litre	10 Lit/litre	-
Apple	22.5 Kg/ha	30 Kg/ha	10 Lit/ha	-
Scab	1.25-1.5	1.5-2	750	-
Grapes	1.25-1.5	1.5-2	750	-
Angular leaf spot	1.25-1.5	1.5-2	750	-
Downy	1.25-1.5	1.5-2	750	-
Mildew	1.25-1.5	1.5-2	750	-
Anthracnose	1.25-1.5	1.5-2	750	-
Guanava	15 Grams	20 Grams	10 Lit/litre	-

Direction of Use :
Many foliar spray and also soil drench, knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer, ASPECT-HTP power sprayer.

CAUTION: WHILE HANDLING THE FUNGICIDE WORKERS AND USERS MUST WEAR FULL PROTECTIVE CLOTHING LIKE LONG SLEEVED SHIRTS, LONG PANTS, CHEMICAL RESISTANT GLOVES, SHOES, GOGGLES, HAT AND MASK.

- PRECAUTIONS:**
1. Avoid contact with foodstuffs, empty foodstuff containers and animals food.
 2. Avoid contact with mouth, eyes, and skin.
 3. Avoid contact with clothing, shoes, and skin.
 4. Avoid contact with the direction of wind.
 5. Avoid contact with the spray mist. Stay in the direction of wind.
 6. Wash thoroughly the contaminated clothes and parts of the body after spraying.
 7. Do not smoke, drink and chew anything while spraying.
 8. Wear full protective clothing while mixing and spraying.
 9. Do not use for spraying on wet surfaces.

First Aid:
1. If swallowed, induce vomiting by licking the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient is unconscious.
2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with copious amount of soap and water.
3. If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
4. If inhaled, remove the patient to fresh air.
5. If on skin, wash with plenty of water.
6. If on clothes, wash with plenty of water.
7. If on shoes, wash with plenty of water.
8. If on hands, wash with plenty of water.
9. If on face, wash with plenty of water.
10. If on hair, wash with plenty of water.
11. If on eyes, wash with plenty of water.
12. If on mouth, wash with plenty of water.
13. If on nose, wash with plenty of water.
14. If on ears, wash with plenty of water.
15. If on neck, wash with plenty of water.
16. If on chest, wash with plenty of water.
17. If on abdomen, wash with plenty of water.
18. If on back, wash with plenty of water.
19. If on legs, wash with plenty of water.
20. If on feet, wash with plenty of water.
21. If on hands, wash with plenty of water.
22. If on face, wash with plenty of water.
23. If on hair, wash with plenty of water.
24. If on clothes, wash with plenty of water.
25. If on shoes, wash with plenty of water.

అగస్త్యా ఎమ్-45

మన్కోజెబ్ 75% డబ్బుల్ ట్రిప్లీ (స్పృశ్యతా సంక్రమణ రోగ నిరోధకం)

Caution: "Not To Be Used On Crops Other Than Specified On This Label/leaflet".
It is a co-ordinated product of Zinc & Manganese Enhance the fungicidal, bactericidal and contact fungicide activity of Mancozeb 75% WP, containing 75% WW active ingredient and balance adjuvants.

Common name of Crop/ Diseases	A.I. (Kg)	Doseage / Ha Formulation (Kg)	Dilution In Water (Lit)	Waiting Period last spray to harvest (In days)
Tomato	1.25-1.5	1.5-2	750	-
Early Blight	1.25-1.5	1.5-2	750	-
Late Blight	1.25-1.5	1.5-2	750	-
Buck Eye Rot	1.25-1.5	1.5-2	750	-
Leaf Spot	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Wheat	1.25-1.5	1.5-2	750	-
Black Rust	1.25-1.5	1.5-2	750	-
Blight	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Downy Mildew	1.25-1.5	1.5-2	750	-
Paddy	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Tip Rot	1.25-1.5	1.5-2	1000	-
Banana	1.25-1.5	1.5-2	1000	-
Sigatoka	1.25-1.5	1.5-2	1000	-
Leaf spot	1.25-1.5	1.5-2	1000	-
Cucumber	22.5 Gram/litre	30 Gram/litre	10 Lit/litre	-
Apple	22.5 Kg/ha	30 Kg/ha	10 Lit/ha	-
Scab	1.25-1.5	1.5-2	750	-
Grapes	1.25-1.5	1.5-2	750	-
Angular leaf spot	1.25-1.5	1.5-2	750	-
Downy	1.25-1.5	1.5-2	750	-
Mildew	1.25-1.5	1.5-2	750	-
Anthracnose	1.25-1.5	1.5-2	750	-
Guanava	15 Grams	20 Grams	10 Lit/litre	-

Direction of Use :
Many foliar spray and also soil drench, knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer, ASPECT-HTP power sprayer.

CAUTION: WHILE HANDLING THE FUNGICIDE WORKERS AND USERS MUST WEAR FULL PROTECTIVE CLOTHING LIKE LONG SLEEVED SHIRTS, LONG PANTS, CHEMICAL RESISTANT GLOVES, SHOES, GOGGLES, HAT AND MASK.

PRECAUTIONS:

1. Avoid contact with foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes, and skin.
3. Avoid contact with clothing, shoes, and skin.
4. Avoid contact with the direction of wind.
5. Avoid contact with the spray mist. Stay in the direction of wind.
6. Wash thoroughly the contaminated clothes and parts of the body after spraying.
7. Do not smoke, drink and chew anything while spraying.
8. Wear full protective clothing while mixing and spraying.
9. Do not use for spraying on wet surfaces.

అగస్త్యా ఎమ్-45

మన్కోజెబ్ 75% డబ్బుల్ ట్రిప్లీ (స్పృశ్యతా సంక్రమణ రోగ నిరోధకం)

Caution: "Not To Be Used On Crops Other Than Specified On This Label/leaflet".
It is a co-ordinated product of Zinc & Manganese Enhance the fungicidal, bactericidal and contact fungicide activity of Mancozeb 75% WP, containing 75% WW active ingredient and balance adjuvants.

Common name of Crop/ Diseases	A.I. (Kg)	Doseage / Ha Formulation (Kg)	Dilution In Water (Lit)	Waiting Period last spray to harvest (In days)
Tomato	1.25-1.5	1.5-2	750	-
Early Blight	1.25-1.5	1.5-2	750	-
Late Blight	1.25-1.5	1.5-2	750	-
Buck Eye Rot	1.25-1.5	1.5-2	750	-
Leaf Spot	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Wheat	1.25-1.5	1.5-2	750	-
Black Rust	1.25-1.5	1.5-2	750	-
Blight	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Downy Mildew	1.25-1.5	1.5-2	750	-
Paddy	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Tip Rot	1.25-1.5	1.5-2	1000	-
Banana	1.25-1.5	1.5-2	1000	-
Sigatoka	1.25-1.5	1.5-2	1000	-
Leaf spot	1.25-1.5	1.5-2	1000	-
Cucumber	22.5 Gram/litre	30 Gram/litre	10 Lit/litre	-
Apple	22.5 Kg/ha	30 Kg/ha	10 Lit/ha	-
Scab	1.25-1.5	1.5-2	750	-
Grapes	1.25-1.5	1.5-2	750	-
Angular leaf spot	1.25-1.5	1.5-2	750	-
Downy	1.25-1.5	1.5-2	750	-
Mildew	1.25-1.5	1.5-2	750	-
Anthracnose	1.25-1.5	1.5-2	750	-
Guanava	15 Grams	20 Grams	10 Lit/litre	-

Direction of Use :
Many foliar spray and also soil drench, knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer, ASPECT-HTP power sprayer.

CAUTION: WHILE HANDLING THE FUNGICIDE WORKERS AND USERS MUST WEAR FULL PROTECTIVE CLOTHING LIKE LONG SLEEVED SHIRTS, LONG PANTS, CHEMICAL RESISTANT GLOVES, SHOES, GOGGLES, HAT AND MASK.

PRECAUTIONS:

1. Avoid contact with foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes, and skin.
3. Avoid contact with clothing, shoes, and skin.
4. Avoid contact with the direction of wind.
5. Avoid contact with the spray mist. Stay in the direction of wind.
6. Wash thoroughly the contaminated clothes and parts of the body after spraying.
7. Do not smoke, drink and chew anything while spraying.
8. Wear full protective clothing while mixing and spraying.
9. Do not use for spraying on wet surfaces.

అగస్త్యా ఎమ్-45

మన్కోజెబ్ 75% డబ్బుల్ ట్రిప్లీ (స్పృశ్యతా సంక్రమణ రోగ నిరోధకం)

Caution: "Not To Be Used On Crops Other Than Specified On This Label/leaflet".
It is a co-ordinated product of Zinc & Manganese Enhance the fungicidal, bactericidal and contact fungicide activity of Mancozeb 75% WP, containing 75% WW active ingredient and balance adjuvants.

Common name of Crop/ Diseases	A.I. (Kg)	Doseage / Ha Formulation (Kg)	Dilution In Water (Lit)	Waiting Period last spray to harvest (In days)
Tomato	1.25-1.5	1.5-2	750	-
Early Blight	1.25-1.5	1.5-2	750	-
Late Blight	1.25-1.5	1.5-2	750	-
Buck Eye Rot	1.25-1.5	1.5-2	750	-
Leaf Spot	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Wheat	1.25-1.5	1.5-2	750	-
Black Rust	1.25-1.5	1.5-2	750	-
Blight	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Downy Mildew	1.25-1.5	1.5-2	750	-
Paddy	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Tip Rot	1.25-1.5	1.5-2	1000	-
Banana	1.25-1.5	1.5-2	1000	-
Sigatoka	1.25-1.5	1.5-2	1000	-
Leaf spot	1.25-1.5	1.5-2	1000	-
Cucumber	22.5 Gram/litre	30 Gram/litre	10 Lit/litre	-
Apple	22.5 Kg/ha	30 Kg/ha	10 Lit/ha	-
Scab	1.25-1.5	1.5-2	750	-
Grapes	1.25-1.5	1.5-2	750	-
Angular leaf spot	1.25-1.5	1.5-2	750	-
Downy	1.25-1.5	1.5-2	750	-
Mildew	1.25-1.5	1.5-2	750	-
Anthracnose	1.25-1.5	1.5-2	750	-
Guanava	15 Grams	20 Grams	10 Lit/litre	-

Direction of Use :
Many foliar spray and also soil drench, knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer, ASPECT-HTP power sprayer.

CAUTION: WHILE HANDLING THE FUNGICIDE WORKERS AND USERS MUST WEAR FULL PROTECTIVE CLOTHING LIKE LONG SLEEVED SHIRTS, LONG PANTS, CHEMICAL RESISTANT GLOVES, SHOES, GOGGLES, HAT AND MASK.

PRECAUTIONS:

1. Avoid contact with foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes, and skin.
3. Avoid contact with clothing, shoes, and skin.
4. Avoid contact with the direction of wind.
5. Avoid contact with the spray mist. Stay in the direction of wind.
6. Wash thoroughly the contaminated clothes and parts of the body after spraying.
7. Do not smoke, drink and chew anything while spraying.
8. Wear full protective clothing while mixing and spraying.
9. Do not use for spraying on wet surfaces.

అగస్త్యా ఎమ్-45

మన్కోజెబ్ 75% డబ్బుల్ ట్రిప్లీ (స్పృశ్యతా సంక్రమణ రోగ నిరోధకం)

Caution: "Not To Be Used On Crops Other Than Specified On This Label/leaflet".
It is a co-ordinated product of Zinc & Manganese Enhance the fungicidal, bactericidal and contact fungicide activity of Mancozeb 75% WP, containing 75% WW active ingredient and balance adjuvants.

Common name of Crop/ Diseases	A.I. (Kg)	Doseage / Ha Formulation (Kg)	Dilution In Water (Lit)	Waiting Period last spray to harvest (In days)
Tomato	1.25-1.5	1.5-2	750	-
Early Blight	1.25-1.5	1.5-2	750	-
Late Blight	1.25-1.5	1.5-2	750	-
Buck Eye Rot	1.25-1.5	1.5-2	750	-
Leaf Spot	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Wheat	1.25-1.5	1.5-2	750	-
Black Rust	1.25-1.5	1.5-2	750	-
Blight	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Downy Mildew	1.25-1.5	1.5-2	750	-
Paddy	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Tip Rot	1.25-1.5	1.5-2	1000	-
Banana	1.25-1.5	1.5-2	1000	-
Sigatoka	1.25-1.5	1.5-2	1000	-
Leaf spot	1.25-1.5	1.5-2	1000	-
Cucumber	22.5 Gram/litre	30 Gram/litre	10 Lit/litre	-
Apple	22.5 Kg/ha	30 Kg/ha	10 Lit/ha	-
Scab	1.25-1.5	1.5-2	750	-
Grapes	1.25-1.5	1.5-2	750	-
Angular leaf spot	1.25-1.5	1.5-2	750	-
Downy	1.25-1.5	1.5-2	750	-
Mildew	1.25-1.5	1.5-2	750	-
Anthracnose	1.25-1.5	1.5-2	750	-
Guanava	15 Grams	20 Grams	10 Lit/litre	-

Direction of Use :
Many foliar spray and also soil drench, knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer, ASPECT-HTP power sprayer.

CAUTION: WHILE HANDLING THE FUNGICIDE WORKERS AND USERS MUST WEAR FULL PROTECTIVE CLOTHING LIKE LONG SLEEVED SHIRTS, LONG PANTS, CHEMICAL RESISTANT GLOVES, SHOES, GOGGLES, HAT AND MASK.

PRECAUTIONS:

1. Avoid contact with foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes, and skin.
3. Avoid contact with clothing, shoes, and skin.
4. Avoid contact with the direction of wind.
5. Avoid contact with the spray mist. Stay in the direction of wind.
6. Wash thoroughly the contaminated clothes and parts of the body after spraying.
7. Do not smoke, drink and chew anything while spraying.
8. Wear full protective clothing while mixing and spraying.
9. Do not use for spraying on wet surfaces.

అగస్త్యా ఎమ్-45

మన్కోజెబ్ 75% డబ్బుల్ ట్రిప్లీ (స్పృశ్యతా సంక్రమణ రోగ నిరోధకం)

Caution: "Not To Be Used On Crops Other Than Specified On This Label/leaflet".
It is a co-ordinated product of Zinc & Manganese Enhance the fungicidal, bactericidal and contact fungicide activity of Mancozeb 75% WP, containing 75% WW active ingredient and balance adjuvants.

Common name of Crop/ Diseases	A.I. (Kg)	Doseage / Ha Formulation (Kg)	Dilution In Water (Lit)	Waiting Period last spray to harvest (In days)
Tomato	1.25-1.5	1.5-2	750	-
Early Blight	1.25-1.5	1.5-2	750	-
Late Blight	1.25-1.5	1.5-2	750	-
Buck Eye Rot	1.25-1.5	1.5-2	750	-
Leaf Spot	1.25-1.5	1.5-2	750	-
Leaf Blight	1.25-1.5	1.5-2	750	-
Wheat	1.25-1.5	1.5-2	750	-
Black Rust	1.25-1.5	1.5-2	750	-
Blight	1.2			

