

AGASTYA GOLD

Bispyribac sodium 10% w/v SC

Systemic Herbicide

Caution : Not To Be Used On Crops Other Than Specified On This Label/Leaflet.
Bispyribac sodium 10% SC is a post emergence herbicides effective against grasses, sedges and broad leaf weeds of rice crop.

Recommendation:	Crop Common name of pests	Dosage/ha	Waiting Period
	A.I. (gm.)	Formula-tion (Ltr.)	Dilution In water Spray to last Harvest (Days)
	20	200	300

Direction of Use: Shake the bottle before use. Measure out requisite quantity of the herbicides and mix it well with small quantity of water. Add the remaining quantity of water specified with thorough agitation.

Application method: Prior to herbicides application the field should be drained. Field should be re-flooded within one day after application and water should be maintained at least for 10 days post application.

Spraying Instruction: Spray uniformly in the direction of wind. Ensure thorough coverage of the weeds.

Time of Application: Nursery/transplanted: The optimum time of the application is 10-12 days after sowing for nursery rice and 10-14 days after planting for transplanted rice when the most of the weeds have already emerged and are of 3-4 leaf stage depending upon soil and climatic factors. Direct seed Rice: For direct seed sowing the time of application is 15-25 days after sowing. Plant Protection Equipment: High volume sprayer w/z. Knapsack sprayer fitted with flat fan hood preffered is recommended.

Precaution: 1. Keep away from food and feedstuff. 2.5 Avoid contact with eyes and skin. 3. Avoid inhalation of the spray mist. 4. Wash thoroughly the contaminated clothes and parts of the body after spraying. 5. Do not smoke, drink eat and chew anything while spraying.

Physio-toxicity: The product is not physio-toxic when used as per field recommendation.

Symptoms of poisoning: Weakness, vomiting, salivation and dizziness. **First Aid:** 1. If swallowed, induce vomiting by tickling the back of throat. Do not induce vomiting if the patient is unconscious. 2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with plenty of saline/clean water for about 10 to 15 minutes. 4. If flushed with plenty of saline/clean water for about 10 to 15 minutes. 4. If inhaled, remove the patient to fresh air, if not breathing, clear the airway and give artificial respiration. Get a medical attention immediately.

Antidote: No specific antidote is known. Treat Symptomatically.

Dispose of used container: 1. Empty containers shall be broken and buried away from habitation. 2. The used containers or packages shall not be left outside to prevent their re-use. 3. The packages of surplus material and waste should be disposed off in a safe manner so as to prevent environmental and water pollution.

Storage Conditions: 1. The packages containing the insecticides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate articles under lock and key depending upon the quantity and nature of the insecticide. 2. The rooms or premises meant for storing the insecticides shall be well built, dry, well-ill, ventilated and of sufficient dimensions to avoid contaminated vapours.

Chemical Composition:

Bispyribac sodium (a.i.)	10.00% w/v
Adjuvant-A: Octadecylamine oxide derivative of Synthetic alcohol	31.90% w/v
Adjuvant-B: Polyoxyethylene Polyoxy Propylene Synthetic alcohol ether	53.20% w/v
Propylene glycol	1.00% w/v
Aluminium Oxide C	1.00% w/v
Silicon KS-506	1.00% w/v
Stabilizer	0.5% w/v
Water	100.00 w/v

AGASTYA GOLD

Bispyribac sodium 10% w/v SC

Systemic Herbicide

Caution : Not To Be Used On Crops Other Than Specified On This Label/Leaflet.
Bispyribac sodium 10% SC is a post emergence herbicides effective against grasses, sedges and broad leaf weeds of rice crop.

Recommendation:	Crop Common name of pests	Dosage/ha	Waiting Period
	A.I. (gm.)	Formula-tion (Ltr.)	Dilution In water Spray to last Harvest (Days)
	20	200	300

Direction of Use: Shake the bottle before use. Measure out requisite quantity of the herbicides and mix it well with small quantity of water. Add the remaining quantity of water specified with thorough agitation.

Application method: Prior to herbicides application the field should be drained. Field should be re-flooded within one day after application and water should be maintained at least for 10 days post application.

Spraying Instruction: Spray uniformly in the direction of wind. Ensure thorough coverage of the weeds.

Time of Application: Nursery/transplanted: The optimum time of the application is 10-12 days after sowing for nursery rice and 10-14 days after planting for transplanted rice when the most of the weeds have already emerged and are of 3-4 leaf stage depending upon soil and climatic factors. Direct seed Rice: For direct seed sowing the time of application is 15-25 days after sowing. Plant Protection Equipment: High volume sprayer w/z. Knapsack sprayer fitted with flat fan hood preffered is recommended.

Precaution: 1. Keep away from food and feedstuff. 2.5 Avoid contact with eyes and skin. 3. Avoid inhalation of the spray mist. 4. Wash thoroughly the contaminated clothes and parts of the body after spraying. 5. Do not smoke, drink eat and chew anything while spraying.

Physio-toxicity: The product is not physio-toxic when used as per field recommendation.

Symptoms of poisoning: Weakness, vomiting, salivation and dizziness. **First Aid:** 1. If swallowed, induce vomiting by tickling the back of throat. Do not induce vomiting if the patient is unconscious. 2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with plenty of saline/clean water for about 10 to 15 minutes. 4. If flushed with plenty of saline/clean water for about 10 to 15 minutes. 4. If inhaled, remove the patient to fresh air, if not breathing, clear the airway and give artificial respiration. Get a medical attention immediately.

Antidote: No specific antidote is known. Treat Symptomatically.

Dispose of used container: 1. Empty containers shall be broken and buried away from habitation. 2. The used containers or packages shall not be left outside to prevent their re-use. 3. The packages of surplus material and waste should be disposed off in a safe manner so as to prevent environmental and water pollution.

Storage Conditions: 1. The packages containing the insecticides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate articles under lock and key depending upon the quantity and nature of the insecticide. 2. The rooms or premises meant for storing the insecticides shall be well built, dry, well-ill, ventilated and of sufficient dimensions to avoid contaminated vapours.

Chemical Composition:

Bispyribac sodium (a.i.)	10.00% w/v
Adjuvant-A: Octadecylamine oxide derivative of Synthetic alcohol	31.90% w/v
Adjuvant-B: Polyoxyethylene Polyoxy Propylene Synthetic alcohol ether	53.20% w/v
Propylene glycol	1.00% w/v
Aluminium Oxide C	1.00% w/v
Silicon KS-506	1.00% w/v
Stabilizer	0.5% w/v
Water	100.00 w/v

AGASTYA GOLD

Bispyribac sodium 10% w/v SC

Systemic Herbicide

Caution : Not To Be Used On Crops Other Than Specified On This Label/Leaflet.
Bispyribac sodium 10% SC is a post emergence herbicides effective against grasses, sedges and broad leaf weeds of rice crop.

Recommendation:	Crop Common name of pests	Dosage/ha	Waiting Period
	A.I. (gm.)	Formula-tion (Ltr.)	Dilution In water Spray to last Harvest (Days)
	20	200	300

Direction of Use: Shake the bottle before use. Measure out requisite quantity of the herbicides and mix it well with small quantity of water. Add the remaining quantity of water specified with thorough agitation.

Application method: Prior to herbicides application the field should be drained. Field should be re-flooded within one day after application and water should be maintained at least for 10 days post application.

Spraying Instruction: Spray uniformly in the direction of wind. Ensure thorough coverage of the weeds.

Time of Application: Nursery/transplanted: The optimum time of the application is 10-12 days after sowing for nursery rice and 10-14 days after planting for transplanted rice when the most of the weeds have already emerged and are of 3-4 leaf stage depending upon soil and climatic factors. Direct seed Rice: For direct seed sowing the time of application is 15-25 days after sowing. Plant Protection Equipment: High volume sprayer w/z. Knapsack sprayer fitted with flat fan hood preffered is recommended.

Precaution: 1. Keep away from food and feedstuff. 2.5 Avoid contact with eyes and skin. 3. Avoid inhalation of the spray mist. 4. Wash thoroughly the contaminated clothes and parts of the body after spraying. 5. Do not smoke, drink eat and chew anything while spraying.

Physio-toxicity: The product is not physio-toxic when used as per field recommendation.

Symptoms of poisoning: Weakness, vomiting, salivation and dizziness. **First Aid:** 1. If swallowed, induce vomiting by tickling the back of throat. Do not induce vomiting if the patient is unconscious. 2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with plenty of saline/clean water for about 10 to 15 minutes. 4. If flushed with plenty of saline/clean water for about 10 to 15 minutes. 4. If inhaled, remove the patient to fresh air, if not breathing, clear the airway and give artificial respiration. Get a medical attention immediately.

Antidote: No specific antidote is known. Treat Symptomatically.

Dispose of used container: 1. Empty containers shall be broken and buried away from habitation. 2. The used containers or packages shall not be left outside to prevent their re-use. 3. The packages of surplus material and waste should be disposed off in a safe manner so as to prevent environmental and water pollution.

Storage Conditions: 1. The packages containing the insecticides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate articles under lock and key depending upon the quantity and nature of the insecticide. 2. The rooms or premises meant for storing the insecticides shall be well built, dry, well-ill, ventilated and of sufficient dimensions to avoid contaminated vapours.

Chemical Composition:

Bispyribac sodium (a.i.)	10.00% w/v
Adjuvant-A: Octadecylamine oxide derivative of Synthetic alcohol	31.90% w/v
Adjuvant-B: Polyoxyethylene Polyoxy Propylene Synthetic alcohol ether	53.20% w/v
Propylene glycol	1.00% w/v
Aluminium Oxide C	1.00% w/v
Silicon KS-506	1.00% w/v
Stabilizer	0.5% w/v
Water	100.00 w/v

AGASTYA GOLD

Bispyribac sodium 10% w/v SC

Systemic Herbicide

Caution : Not To Be Used On Crops Other Than Specified On This Label/Leaflet.
Bispyribac sodium 10% SC is a post emergence herbicides effective against grasses, sedges and broad leaf weeds of rice crop.

Recommendation:	Crop Common name of pests	Dosage/ha	Waiting Period
	A.I. (gm.)	Formula-tion (Ltr.)	Dilution In water Spray to last Harvest (Days)
	20	200	300

Direction of Use: Shake the bottle before use. Measure out requisite quantity of the herbicides and mix it well with small quantity of water. Add the remaining quantity of water specified with thorough agitation.

Application method: Prior to herbicides application the field should be drained. Field should be re-flooded within one day after application and water should be maintained at least for 10 days post application.

Spraying Instruction: Spray uniformly in the direction of wind. Ensure thorough coverage of the weeds.

Time of Application: Nursery/transplanted: The optimum time of the application is 10-12 days after sowing for nursery rice and 10-14 days after planting for transplanted rice when the most of the weeds have already emerged and are of 3-4 leaf stage depending upon soil and climatic factors. Direct seed Rice: For direct seed sowing the time of application is 15-25 days after sowing. Plant Protection Equipment: High volume sprayer w/z. Knapsack sprayer fitted with flat fan hood preffered is recommended.

Precaution: 1. Keep away from food and feedstuff. 2.5 Avoid contact with eyes and skin. 3. Avoid inhalation of the spray mist. 4. Wash thoroughly the contaminated clothes and parts of the body after spraying. 5. Do not smoke, drink eat and chew anything while spraying.

Physio-toxicity: The product is not physio-toxic when used as per field recommendation.

Symptoms of poisoning: Weakness, vomiting, salivation and dizziness. **First Aid:** 1. If swallowed, induce vomiting by tickling the back of throat. Do not induce vomiting if the patient is unconscious. 2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with plenty of saline/clean water for about 10 to 15 minutes. 4. If flushed with plenty of saline/clean water for about 10 to 15 minutes. 4. If inhaled, remove the patient to fresh air, if not breathing, clear the airway and give artificial respiration. Get a medical attention immediately.

Antidote: No specific antidote is known. Treat Symptomatically.

Dispose of used container: 1. Empty containers shall be broken and buried away from habitation. 2. The used containers or packages shall not be left outside to prevent their re-use. 3. The packages of surplus material and waste should be disposed off in a safe manner so as to prevent environmental and water pollution.


Storage Conditions: 1. The packages containing the insecticides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate articles under lock and key depending upon the quantity and nature of the insecticide. 2. The rooms or premises meant for storing the insecticides shall be well built, dry, well-ill, ventilated and of sufficient dimensions to avoid contaminated vapours.

Chemical Composition:

Bispyribac sodium (a.i.)	10.00% w/v
Adjuvant-A: Octadecylamine oxide derivative of Synthetic alcohol	31.90% w/v
Adjuvant-B: Polyoxyethylene Polyoxy Propylene Synthetic alcohol ether	53.20% w/v
Propylene glycol	1.00% w/v
Aluminium Oxide C	1.00% w/v
Silicon KS-506	1.00% w/v
Stabilizer	0.5% w/v
Water	100.00 w/v

INSTRUCTIONS FOR USE

Keep in cool, dry place away from heat and open flame.
Do not use on crops other than specified on this label.
KEEP OUT OF THE REACH OF CHILDREN.
Do not use on crops other than specified on this label.
Do not use on crops other than specified on this label.



READ THE ENTIRE LABEL AND LABEL BEFORE USE.
AGASTYA GOLD is a systemic herbicide.
For agricultural use only.

Manufactured and Marketed by:
AGASTYA AGRO LIMITED
An ISO Certified Company
UNIT-1, Survey No. 47, Muraraj Pally - 501 401
Medical Branch, Medical - Malajpet Dist. Telangana State.
Mfg. Lic. No. 20775-MIL-2017
OC Lic. No. 2207-997-991 Dt. 06-07-1997

UNIT-4, H.No. 245, St. No. 54/1 & 55,
Footcoper (V), Chouhanpet (M),
Yadadri-Bhuvanagiri Dist. 508 252 TS.
Mfg. Lic. No. 22175-MIL-2015
OC Lic. No. 2207-997-991 Dt. 06-07-1997

Customer Care Manager: Mr. 94464578988
E-mail: agastyaagro@gmail.com

Regd. No. DR-100015/2015-Bispyribac Sodium (SC) (405)-56



